

БЛАНК ОТВЕТОВ

[illegible]

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100



Возраст

--	--	--

Военнослужащий ☐Пенсионер ☐

Иное ☐

e-mail

Телефон

Подпись участника строго внутри окна

☐

Будьте аккуратны. Случайный штрих внутри квадрата может быть воспринят как метка

Варианты ответов

[illegible][illegible]

Figure 1 is a schematic representation of the experimental design. It shows two groups of subjects: 'Control' and 'Patients'. Each group has a 'Pre' measurement and a 'Post' measurement. The 'Control' group is represented by a solid black rectangle, and the 'Patients' group is represented by a white rectangle with a black border. The 'Pre' measurement is taken before the intervention, and the 'Post' measurement is taken after the intervention. The 'Patients' group shows a significant decrease in the 'Post' measurement compared to the 'Control' group, indicated by a downward arrow and a p-value of $p < 0.05$.

☐

Подпись ответственного организатора
строго внутри окна